

Strongly modular lattices with long shadow.

Gabriele Nebe

Abteilung Reine Mathematik, Universität Ulm, 89069 Ulm, Germany

nebe@mathematik.uni-ulm.de

February 1, 2008

ABSTRACT:* We classify strongly modular lattices with longest and second longest possible shadow.

1 Introduction

To an integral lattice L in the euclidean space $(\mathbb{R}^n, (\cdot, \cdot))$, one associates the set of **characteristic vectors** $v \in \mathbb{R}^n$ with $(v, x) \equiv (x, x) \pmod{2\mathbb{Z}}$ for all $x \in L$. They form a coset modulo $2L^*$, where

$$L^* = \{v \in \mathbb{R}^n \mid (v, x) \in \mathbb{Z} \ \forall x \in L\}$$

is the **dual lattice** of L . Recall that L is called **integral**, if $L \subset L^*$ and **unimodular**, if $L = L^*$. For a unimodular lattice, the square length of a characteristic vector is congruent to n modulo 8 and there is always a characteristic vector of square length $\leq n$. In [1] Elkies characterized the standard lattice $\mathbb{Z}^n$ as the unique unimodular lattice of dimension n , for which all characteristic vectors have square length $\geq n$. [2] gives the short list of unimodular lattices L with $\min(L) \geq 2$ such that all characteristic vectors of L have length $\geq n - 8$. The largest dimension n is 23 and in dimension 23 this lattice is the **shorter Leech lattice** O_{23} of minimum 3. In this paper, these theorems are generalized to certain strongly modular lattices. Following [7] and [8], an integral lattice L is called **N -modular**, if L is isometric to its rescaled dual lattice $\sqrt{N}L^*$. A N -modular lattice L is called **strongly N -modular**, if L is isometric to all rescaled partial dual lattices $\sqrt{m}L^{*,m}$, for all exact divisors m of N , where

$$L^{*,m} := L^* \cap \frac{1}{m}L.$$

The simplest strongly N -modular lattice is

$$C_N := \perp_{d|N} \sqrt{d}\mathbb{Z}$$

of dimension $\sigma_0(N) := \sum_{d|N} 1$ the number of divisors of N . The lattice C_N plays the role of $\mathbb{Z} = C_1$ for square free $N > 1$.

*MSC 11H31, 11H50

With the help of modular forms Quebbemann [8] shows that for

$$N \in \mathcal{L} := \{1, 2, 3, 5, 6, 7, 11, 14, 15, 23\}$$

(which is the set of all positive integers N such that the sum of divisors

$$\sigma_1(N) := \sum_{d|N} d$$

divides 24), the minimum of an even strongly N -modular lattice L of dimension n satisfies

$$\min(L) \leq 2 + 2 \lfloor \frac{n \sigma_1(N)}{24 \sigma_0(N)} \rfloor.$$

Strongly modular lattices meeting this bound are called **extremal**. Whereas Quebbemann restricts to even lattices, [9] shows that the same bound also holds for odd strongly modular lattices, where there is one exceptional dimension $n = \sigma_0(N)(\frac{24}{\sigma_1(N)} - 1)$, where the bound on the minimum is 3 (and not 2). In this dimension, there is a unique lattice $S^{(N)}$ of minimum 3. For $N = 1$, this is again the shorter Leech lattice O_{23} . The main tool to get the bound for odd lattices is the **shadow**

$$S(L) := \{ \frac{v}{2} \mid v \text{ is a characteristic vector of } L \}.$$

If L is even, then $S(L) = L^*$ and if L is odd, $S(L) = L_0^* - L^*$, where

$$L_0 := \{v \in L \mid (v, v) \in 2\mathbb{Z}\}$$

is the even sublattice of L .

The main result of this paper is Theorem 3. It is shown that for a strongly N -modular lattice L that is rationally equivalent to C_N^k , the minimum

$$\min_0(S(L)) := \min\{(v, v) \mid v \in S(L)\}$$

equals

$$M^{(N)}(m, k) := \begin{cases} \frac{1}{N}(k \frac{\sigma_1(N)}{4} - 2m) & \text{if } N \text{ is odd} \\ \frac{1}{N}(k \frac{\sigma_1(N/2)}{2} - m) & \text{if } N \text{ is even} \end{cases}$$

for some $m \in \mathbb{Z}_{\geq 0}$. If $\min_0(S(L)) = M^{(N)}(0, k)$, then $L \cong C_N^k$. For the next smaller possible minimum $\min_0(S(L)) = M^{(N)}(1, k)$ one gets that $L \cong C_N^l \perp L'$, where $\min(L') > 1$ and $\dim(L') \leq \sigma_0(N)(s(N) - 1)$ for odd N resp. $\dim(L') \leq \sigma_0(N)s(N)$ for even N . The lattices L' of maximal possible dimensions have minimum 3 and are uniquely determined: $L' = S^{(N)}$, if N is odd and $L' = O^{(N)}$ (the “odd analogue” of the unique extremal strongly N -modular lattice of dimension $\sigma_0(N)s(N)$) if N is even (see [9, Table 1]).

The main tool to prove this theorem are the formulas for the theta series of a strongly N -modular lattice L and of its shadow $S(L)$ developed in [9]. Therefore we briefly repeat these formulas in the next section.

2 Theta series

For a subset $S \subset \mathbb{R}^n$, which is a finite union of cosets of an integral lattice we put its **theta series**

$$\Theta_S(z) := \sum_{v \in S} q^{(v,v)}, \quad q = \exp(\pi iz).$$

The theta series of strongly N -modular lattices are modular forms for a certain discrete subgroup Γ_N of $SL_2(\mathbb{R})$ (see [9]). Fix $N \in \mathcal{L}$ and put

$$g_1^{(N)}(z) := \Theta_{C_N}(z) = 1 + 2q + 2\text{ev}(N)q^2 + \dots$$

where

$$\text{ev}(N) := \begin{cases} 1 & \text{if } N \text{ is even} \\ 0 & \text{if } N \text{ is odd} \end{cases}.$$

Let η be the Dedekind eta-function

$$\eta(z) := q^{\frac{1}{12}} \prod_{m=1}^{\infty} (1 - q^{2m}), \quad q = \exp(\pi iz).$$

and put

$$\eta^{(N)}(z) := \prod_{d|N} \eta(dz).$$

If N is odd define

$$g_2^{(N)}(z) := \left(\frac{\eta^{(N)}(z/2)\eta^{(N)}(2z)}{\eta^{(N)}(z)^2} \right)^s$$

and if N is even then

$$g_2^{(N)}(z) := \left(\frac{\eta^{(N/2)}(z/2)\eta^{(N/2)}(4z)}{\eta^{(N/2)}(z)\eta^{(N/2)}(2z)} \right)^s.$$

Then $g_2^{(N)}$ generates the field of modular functions of Γ_N . It is a power series in q starting with

$$g_2^{(N)}(z) = q - sq^2 + \dots$$

Theorem 1 ([9, Theorem 9, Corollary 3]) *Let $N \in \mathcal{L}$ and L be a strongly N -modular lattice that is rational equivalent to C_N^k . Define $l_N := \frac{1}{8}\sigma_1(N)$, if N is odd and $l_N := \frac{1}{6}\sigma_1(N)$, if N is even. Then*

$$\Theta_L(z) = g_1^{(N)}(z)^k \sum_{i=0}^{\lfloor kl_N \rfloor} c_i g_2^{(N)}(z)^i$$

for $c_i \in \mathbb{R}$. The theta series of the rescaled shadow $S := \sqrt{N}S(L)$ of L is

$$\Theta_S(z) = s_1^{(N)}(z)^k \sum_{i=0}^{\lfloor kl_N \rfloor} c_i s_2^{(N)}(z)^i$$

where $s_1^{(N)}$ and $s_2^{(N)}$ are the corresponding “shadows” of $g_1^{(N)}$ and $g_2^{(N)}$.

For odd N

$$s_1^{(N)}(z) = 2^{\sigma_0(N)} \frac{\eta^{(N)}(2z)^2}{\eta^{(N)}(z)}$$

and

$$s_2^{(N)}(z) = -2^{-s(N)\sigma_0(N)/2} \left(\frac{\eta^{(N)}(z)}{\eta^{(N)}(2z)} \right)^{s(N)}$$

For $N = 2$ one has

$$s_1^{(2)}(z) = \frac{2\eta(z)^5\eta(4z)^2}{\eta(z/2)^2\eta(2z)^3}$$

and

$$s_2^{(2)}(z) = -\frac{1}{16} \left(\frac{\eta(z/2)\eta(2z)^2}{\eta(z)^2\eta(4z)} \right)^8$$

which yields $s_1^{(N)}$ and $s_2^{(N)}$ for $N = 6, 14$ as

$$s_1^{(N)} = s_1^{(2)}(z)s_1^{(2)}\left(\frac{N}{2}z\right)$$

and

$$s_2^{(N)} = (s_2^{(2)}(z)s_2^{(2)}\left(\frac{N}{2}z\right))^{s(N)/s(2)}.$$

If N is odd, then $s_1^{(N)}$ starts with $q^{\sigma_1(N)/4}$ and $s_2^{(N)}$ starts with q^{-2} . If N is even, then $s_1^{(N)}$ starts with $q^{\sigma_1(N/2)/2}$ and $s_2^{(N)}$ starts with q^{-1} .

3 Strongly modular lattices with long shadow.

Proposition 2 *Let $N \in \mathbb{N}$ be square free and let L be a strongly N -modular lattice. If L contains a vector of length 1, then L has an orthogonal summand C_N .*

Proof. Since L is an integral lattice that contains a vector of length 1, the unimodular lattice $\mathbb{Z}$ is an orthogonal summand of L . Hence $L = \mathbb{Z} \perp L'$. If d is a divisor of N , then

$$L \cong \sqrt{d}L^{*,d} = \sqrt{d}\mathbb{Z} \perp \sqrt{d}(L')^{*,d}$$

by assumption. Hence L contains an orthogonal summand $\sqrt{d}\mathbb{Z}$ for all divisors d of N and therefore C_N is an orthogonal summand of L . $\square$

Theorem 3 (see [2] for $N = 1$) *Let $N \in \mathcal{L}$ and L be a strongly N -modular lattice that is rational equivalent to C_N^k . Let $M^{(N)}(m, k)$ be as defined in the introduction.*

(i) $\min_0(S(L)) = M^{(N)}(m, k)$ for some $m \in \mathbb{Z}_{\geq 0}$.

(ii) If $\min_0(S(L)) = M^{(N)}(0, k)$ then $L \cong C_N^k$.

(iii) If $\min_0(S(L)) = M^{(N)}(m, k)$ then $L \cong C_N^a \perp L'$, where L' is a strongly N -modular lattice rational equivalent to C_N^{k-a} with $\min(L') \geq 2$ and $\min_0(S(L')) = M^{(N)}(m, k - a)$.

(iv) If $\min_0(S(L)) = M^{(N)}(m, 1)$ and $\min(L) \geq 2$, then the number of vectors of length 2 in L is

$$2k(s(N) + \text{ev}(N) - (k + 1)).$$

In particular $k \leq k_{\max}(N)$ with

$$k_{\max}(N) = s(N) - 1 + \text{ev}(N)$$

and if $k = k_{\max}(N)$, then $\min(L) \geq 3$.

Proof. (i) follows immediately from Theorem 1.

(ii) In this case the theta series of L is g_1^k . In particular L contains $2k$ vectors of norm 1. Applying Proposition 2 one finds that $L \cong C_N$.

(iii) Follows from Proposition 2 and Theorem 1.

(iv) Since $\min(L) > 1$, $\Theta_L = g_1^k - 2kg_1^k g_2$. Explicit calculations give the number of norm-2-vectors in L . $\square$

The following table gives the maximal dimension $n_{\max}(N) = \sigma_0(N)k_{\max}(N)$ of a lattice in Theorem 3 (iv).

N	1	2	3	5	6	7	11	14	15	23
$\sigma_1(N)$	1	3	4	6	12	8	12	24	24	24
$k_{\max}(N)$	23	8	5	3	2	2	1	1	0	0
$n_{\max}(N)$	23	16	10	6	8	4	2	4	0	0

The lattices L with $\min_0(S(L)) = M(1, k)$ are listed in an appendix. These are only finitely many since k is bounded by $k_{\max}$. In general it is an open problem whether for all m , there are only finitely many strongly N -modular lattices L rational equivalent to C_N^k for some k and of minimum $\min(L) > 1$ such that $\min_0(S(L)) = M(m, k)$. For $N = 1$, Gaultier [3] proved that $k \leq 2907$ for $m = 2$ and $k \leq 8388630$ for $m = 3$.

Theorem 4 (cf. [2] for $N = 1$) Let $N \in \mathcal{L}$ be odd and $k \in \mathbb{N}$ such that

$$\frac{8}{\sigma_1(N)} \leq k \leq k_{\max}(N) = \frac{24}{\sigma_1(N)} - 1.$$

Then there is a unique strongly N -modular lattice $L := L_k(N)$ that is rational equivalent to C_N^k such that $\min(L) > 1$ and $\min_0(S(L)) = M^{(N)}(1, k)$, except for $N = 1$, where there is no such lattice in dimension 9, 10, 11, 13 and there are two lattices in dimension 18 and 20 (see [2]). If $k = k_{\max}(N)$, then L is the shorter lattice $L = S^{(N)}$ described in [9, Table 1] and $\min(L) = 3$.

Proof. For $N = 15$ and $N = 23$ there is nothing to show since $k_{\max}(N) = 0$. The case $N = 1$ is already shown in [2]. It remains to consider $N \in \{3, 5, 7, 11\}$. Since N is a prime, there are only 2 genera of strongly modular lattices, one consisting of even lattices and one of odd lattices. With a short MAGMA program using Kneser's neighboring method, one obtains a list of all lattices in the relevant genus. In all cases there is a unique lattice with the right number of vectors of length 2. Gram matrices of these lattices are given in the appendix. $\square$

Remark 5 For $N = 1$ and dimension $n = 9, 10, 11$ the theta series of the hypothetical shadow has non integral resp. odd coefficients, so there is no lattice $L_n(1)$.

Theorem 6 Let $N \in \mathcal{L}$ be even and $k \in \mathbb{N}$ such that

$$\frac{2}{\sigma_1(N/2)} \leq k \leq k_{\max}(N) = \frac{24}{\sigma_1(N)}.$$

If $(k, N) \neq (3, 2)$ then there are strongly N -modular lattices $L := L_k(N)$ that are rational equivalent to C_N^k such that $\min(L) > 1$ and $\min_0(S(L)) = M^{(N)}(1, k)$. If $k = k_{\max}(N)$, then $L_k(N)$ is unique. It is the odd lattice $L = O^{(N)}$ described in [9, Table 1] and $\min(L) = 3$.

Remark 7 For $N = 2$ and $k = 3$ the corresponding shadow modular form has non integral coefficients, so there is no lattice $L_3(2)$.

Remark 8 All odd lattices $L_k(N)$ in Theorem 6 lie in the genus of C_N^k .

4 Appendix: The lattices $L_k(N)$.

The lattices $L_k(1)$:

The lattices $L_k(1)$ are already listed in [2]. They are uniquely determined by their root-sublattices R_k and given in the following table:

k	8	12	14	15	16	17	18	19	20	21	22	23
R_k	E_8	D_{12}	E_7^2	A_{15}	D_8^2	$A_{11}E_6$	D_6^3, A_9^2	$A_7^2D_5$	D_4^5, A_5^4	A_3^7	A_1^{22}	0

The lattices $L_k(N)$ for $N > 1$ odd:

$$L_2(3): \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \perp \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \cong A_2 \perp A_2, \text{ Automorphism group: } D_{12} \wr C_2.$$

$$L_3(3): \begin{pmatrix} 211111 \\ 121111 \\ 112111 \\ 111300 \\ 111030 \\ 111003 \end{pmatrix} \text{ Automorphism group: order 1152.}$$

$$L_4(3): \begin{pmatrix} 20 & 0 & 0 & -1 & -1 & 0 & 1 \\ 02 & 0 & 0 & 1 & 1 & 1 & 0 \\ 00 & 2 & 0 & 0 & -1 & -1 & -1 \\ 00 & 0 & 2 & -1 & 0 & -1 & -1 \\ -11 & 0 & -1 & 3 & 1 & 1 & 0 \\ -11 & -1 & 0 & 1 & 3 & 1 & 0 \\ 01 & -1 & -1 & 1 & 1 & 3 & 1 \\ 10 & -1 & -1 & 0 & 0 & 1 & 3 \end{pmatrix} \text{ Automorphism group: order 6144.}$$

$$L_5(3): \begin{pmatrix} 3 & -1 & -1 & -1 & 0 & -1 & -1 & 1 & 1 & 0 \\ -1 & 3 & 1 & -1 & -1 & 1 & -1 & 1 & 1 & 0 \\ -1 & 1 & 3 & 1 & 1 & 1 & -1 & 1 & 0 & -1 \\ -1 & -1 & 1 & 3 & 1 & 0 & 1 & 1 & -1 & -1 \\ 0 & -1 & 1 & 1 & 3 & 1 & 0 & -1 & -1 & 0 \\ -1 & 1 & 1 & 0 & 1 & 3 & -1 & 0 & -1 & -1 \\ -1 & -1 & -1 & 1 & 0 & -1 & 3 & 0 & -1 & 1 \\ -1 & 1 & 1 & 1 & -1 & 0 & 0 & 3 & 1 & 0 \\ 1 & 1 & 0 & -1 & -1 & -1 & 1 & 1 & 3 & 1 \\ 0 & 0 & -1 & -1 & 0 & -1 & 1 & 0 & 1 & 3 \end{pmatrix} \text{ Automorphism group: } \pm U_4(2).2 \text{ of order 103680}$$

$$L_2(5): \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix} \perp \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix}, \text{ Automorphism group: } (\pm C_2) \wr C_2 \text{ of order 32.}$$

$$L_3(5): \begin{pmatrix} 3 & -1 & 1 & -1 & 10 \\ -1 & 3 & -1 & 0 & 11 \\ 1 & -1 & 3 & 1 & 01 \\ -1 & 0 & 1 & 3 & -11 \\ 1 & 1 & 0 & -1 & 31 \\ 0 & 1 & 1 & 1 & 13 \end{pmatrix}. \text{ Automorphism group: } \pm S_5 \text{ of order 240.}$$

$$L_1(7): \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}. \text{ Automorphism group: } \pm C_2.$$

$$L_2(7): \begin{pmatrix} 3 & -11 & 0 \\ -1 & 30 & 1 \\ 1 & 03 & 1 \\ 0 & 11 & 3 \end{pmatrix}. \text{ Automorphism group: order 16.}$$

$$L_1(11): \begin{pmatrix} 2 & 1 \\ 1 & 6 \end{pmatrix}. \text{ Automorphism group: } \pm C_2.$$

The lattices $L_k(N)$ for N even:

For $N = 2$ there is only one genus of odd lattices to be considered. Also for $N = 14$ there is only one odd genus for each k , since 2 is a square modulo 7. For $N = 6$, there are 2 such genera, since $L := (\sqrt{2}\mathbb{Z})^2 \perp (\sqrt{3}\mathbb{Z})^2$ is not in the genus of C_6 . The genus of L contains no strongly modular lattices. The genus of $L \perp C_6$ contains 3 lattices with minimum 3, none of which is strongly modular.

$$L_2(2): L_2(2) = D_4 \text{ with automorphism group } W(F_4) \text{ of order 1152.}$$

$$L_4(2) : \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & -1 & -1 & 1 \\ 0 & 2 & 0 & 0 & 0 & 1 & 1 & -1 \\ 0 & 0 & 2 & -1 & 1 & 1 & 1 & -1 \\ 0 & 0 & -1 & 2 & -1 & -1 & 0 & 1 \\ 0 & 0 & 1 & -1 & 2 & 1 & 0 & -1 \\ -1 & 1 & 1 & -1 & 1 & 3 & 2 & -2 \\ -1 & 1 & 1 & 0 & 0 & 2 & 3 & -1 \\ 1 & -1 & -1 & 1 & -1 & -2 & -1 & 3 \end{pmatrix} . \quad \begin{array}{l} \text{The root sublattice is } D_4 \perp A_1^4 \\ \text{and the automorphism group of } L_4(2) \text{ is} \\ W(F_4) \times (C_2^4 : D_8) \text{ of order } 147456. \end{array}$$

$$L_5(2) : \begin{pmatrix} 2 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 2 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 2 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 3 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 1 & 3 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 1 & 1 & 3 & 1 \\ 1 & 1 & 1 & 0 & 1 & 1 & 1 & 1 & 3 \end{pmatrix} . \quad \begin{array}{l} \text{The root sublattice is } A_5 \\ \text{and the automorphism group of } L_5(2) \text{ is} \\ \pm S_6 \times S_6 \text{ of order } 1036800. \end{array}$$

$L_6(2)$: There are two such lattices:

$$L_{6a}(2) : \begin{pmatrix} 2 & 0 & 1 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & -1 & -1 \\ 0 & 2 & 0 & 0 & -1 & 1 & 1 & 1 & -1 & 1 & 0 & -1 \\ 1 & 0 & 2 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & -1 & -1 \\ 1 & 0 & 1 & 2 & 0 & 0 & 1 & 1 & 1 & 1 & -1 & -1 \\ 0 & -1 & 0 & 0 & 2 & -1 & 0 & -1 & 1 & -1 & 1 & 1 \\ 0 & 1 & 0 & 0 & -1 & 2 & 0 & 1 & -1 & 1 & 0 & -1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 3 & 1 & 0 & 1 & 0 & -1 \\ 1 & 1 & 1 & 1 & -1 & 1 & 1 & 3 & 0 & 1 & -1 & -1 \\ 1 & -1 & 1 & 1 & 1 & -1 & 0 & 0 & 3 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & -1 & 1 & 1 & 1 & 0 & 3 & -1 & -1 \\ -1 & 0 & -1 & -1 & 1 & 0 & 0 & -1 & 0 & -1 & 3 & 1 \\ -1 & -1 & -1 & -1 & 1 & -1 & -1 & -1 & 0 & -1 & 1 & 3 \end{pmatrix}, \text{ and } L_{6b}(2) : \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 2 & 0 & 0 & 1 & -1 & 0 & 1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & -1 & 1 & 0 & -1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & -1 & 0 & 1 & 1 \\ 1 & 0 & 1 & -1 & 0 & 3 & 0 & 0 & 1 & -1 & -1 & -1 \\ 1 & 0 & -1 & 1 & 0 & 0 & 3 & -1 & -2 & 2 & 2 & 2 \\ -1 & 0 & 1 & 0 & -1 & 0 & -1 & 3 & 1 & -2 & -2 & -2 \\ -1 & 0 & 1 & -1 & 0 & 1 & -2 & 1 & 3 & -2 & -2 & -2 \\ 1 & 1 & -1 & 0 & 1 & 1 & -1 & 2 & -2 & -2 & 4 & 3 \\ 1 & 1 & -1 & 0 & 1 & 1 & -1 & 2 & -2 & -2 & 3 & 4 \end{pmatrix}$$

with automorphism group of order $2^{15}3^4$ resp. $2^{21}3$.

$$L_7(2) : \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & -1 & 0 \\ 0 & 2 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 2 & 0 & 1 & 1 & 0 & 0 & -1 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & -1 & 0 & -1 & -1 & 0 & 0 & -1 & -1 & 1 & 0 \\ 0 & 1 & -1 & 1 & 3 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 1 & 3 & 1 & 1 & 0 & 1 & 1 & 0 & -1 & 0 \\ 1 & 1 & 0 & -1 & 1 & 1 & 3 & 2 & 0 & 0 & 0 & 0 & -1 & 1 \\ 1 & 1 & 0 & -1 & 1 & 1 & 2 & 3 & 1 & 1 & 1 & 1 & -1 & 0 \\ 1 & 0 & -1 & 0 & 0 & 0 & 0 & 1 & 3 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 & 3 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 & 0 & 1 & 0 & 1 & 3 & 0 & -1 & -1 \\ 0 & 0 & -1 & -1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 3 & -1 & -1 \\ -1 & 0 & 0 & 1 & 0 & -1 & -1 & -1 & 0 & 0 & -1 & -1 & 3 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & -1 & -1 & 1 & 3 \end{pmatrix} \quad \text{Automorphism group of order } 2752512.$$

$L_8(2)$: $L_8(2)$ is the odd version of the Barnes Wall lattice BW_{16} (see [6]). It is unique by [9, Theorem 8].

$$L_1(6): \begin{pmatrix} 2 & -1 & -1 & 0 \\ -1 & 3 & 0 & -1 \\ -1 & 0 & 3 & -1 \\ 0 & -1 & -1 & 4 \end{pmatrix}. \text{ Automorphism group } C_2^4.$$

$$L_2(6): \begin{pmatrix} 3 & 1 & 0 & 0 & 0 & 1 & 0 & -1 \\ 1 & 3 & 0 & 0 & -1 & 0 & -1 & 0 \\ 0 & 0 & 3 & 1 & 0 & -1 & 0 & -1 \\ 0 & 0 & 1 & 3 & 1 & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 & 3 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 3 & 0 & 0 \\ 0 & -1 & 0 & -1 & 0 & 0 & 3 & -1 \\ -1 & 0 & -1 & 0 & 0 & 0 & -1 & 3 \end{pmatrix} \text{ Automorphism group } SL_2(3).2^2 \text{ of order 96.}$$

$$L_1(14): \text{ Gram matrix } \begin{pmatrix} 3 & 1 \\ 1 & 5 \end{pmatrix} \perp \begin{pmatrix} 3 & 1 \\ 1 & 5 \end{pmatrix}. \text{ Automorphism group } D_8.$$

References

- [1] *N.D. Elkies*: A characterization of the $\mathbb{Z}^n$ lattice. Math. Res. Lett. 2 (1995), no. 3, 321-326.
- [2] *N.D. Elkies*: Lattices and codes with long shadows. Math. Res. Lett. 2 (1995), no. 5, 643-651
- [3] *M. Gaultier*: Lattices without short characteristic vectors. Math. Res. Lett. 5 (1998), no. 3, 353-362.
- [4] *C. L. Mallows, A. M. Odlysko, N. J. A. Sloane*: Upper bounds for modular forms, lattices and codes. J. Alg. **36** (1975), 68-76.
- [5] *T. Miyake*: Modular Forms. Springer (1989)
- [6] *G. Nebe, N.J.A. Sloane*: A database of lattices. www.research.att.com/~njas/lattices
- [7] *H.-G. Quebbemann*: Modular lattices in euclidean spaces. J. Number Th. **54** (1995), 190-202.
- [8] *H.-G. Quebbemann*: Atkin-Lehner eigenforms and strongly modular lattices. L'Ens. Math. **43** (1997), 55-65.
- [9] *E.M. Rains, N.J.A. Sloane*: The shadow theory of modular and unimodular lattices. J. Number Th. **73** (1998), 359-389.